

REPORT

Name	: Mrs. CH SUMALATHA	Sample ID	: 24854229
Age/Gender	: 44 Years/Female	Reg. No	: 0312309270051
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 27-Sep-2023 02:31 PM
Primary Sample	: Whole Blood	Received On	: 27-Sep-2023 04:42 PM
Sample Tested In	: Whole Blood EDTA	Reported On	: 27-Sep-2023 05:11 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

HAEMATOLOGY

Test Name	Results	Units	Ref. Range	Method
Haemoglobin (Hb)	7.7	g/dL	12-15	Cynmeth Method

Result rechecked and verified for abnormal cases

*** End Of Report ***

Laboratory is NABL Accredited



*TESTS CONDUCTED @ CENTRAL LAB, HYDERABAD

terms and conditions overleaf. Partial Reproduction of this report is not Permitted

Swarnabala . M
DR.SWARNA BALA
MD PATHOLOGY

REPORT

Name	: Mrs. CH SUMALATHA	Sample ID	: 24854231
Age/Gender	: 44 Years/Female	Reg. No	: 0312309270051
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 27-Sep-2023 02:31 PM
Primary Sample	: Whole Blood	Received On	: 27-Sep-2023 04:50 PM
Sample Tested In	: Serum	Reported On	: 27-Sep-2023 06:41 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Ref. Range	Method
TSH -Thyroid Stimulating Hormone	9.68	μ U/mL	0.35-5.5	CLIA

Pregnancy & Cord Blood

TSH (Thyroid Stimulating Hormone (μ U/mL))	
First Trimester	: 0.24-2.99
Second Trimester	: 0.46-2.95
Third Trimester	: 0.43-2.78
Cord Blood	: 2.3-13.2

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

Correlate Clinically.

Result rechecked and verified for abnormal cases

Laboratory is NABL Accredited

*** End Of Report ***



Dr. Vaishnavi
DR. VAISHNAVI
MD BIOCHEMISTRY