

Sagepath Labs Pvt. Ltd.

Registered Office:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

	-	^	D '	-	
R	-	()	к.		
	_	~	•		

KEPOKI							
Name	: Mrs. KAVITHA	Sample ID	: 24753475				
Age/Gender	: 31 Years/Female	Reg. No	: 0312311140050				
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172				
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 14-Nov-2023 09:11 PM				
Primary Sample	: Whole Blood	Received On	: 15-Nov-2023 11:34 AM				
Sample Tested In	: Serum	Reported On	: 15-Nov-2023 11:35 AM				
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report				

	IMMUNOLOGY & SEROLOGY			
Test Name	Results	Units	Ref. Range	Method
HIV 123 Profile				
HIV (1& 2) Antibody	3.49	S/Co	< 1.00 : Negative > 1.00 : Positive	ELISA
HIV 1 &2 Ab-Chromatography				
HIV - I Results	Reactive		Non Reactive	Immuno Chromatography
HIV-II Results	Non Reactive		Non Reactive	Immunochromatography
HIV-Immunofilteration				
HIV 1	Reactive		Non-reactive	Immunofiltration
HIV 2	Non Reactive		Non Reactive	Immunofiltration
Final Report.				
Final Report-	Tested sample Positive for Hiv-1 antibody			

Interpretation

• Non Reactive result implies that antibodies to HIV 1 / 2 have not been detected in the sample. This means the patient has either not been exposed to HIV 1 / 2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1 / 2.

• Pre and Post test counseling to be done by the concerned referring doctor. The sensitivity and specificity of this test has been determined by National HIV Reference Centers of Govt. of India and WHO collaborating Centers, using various other test panels."

• Reactive samples by ELISA Method are confirmed by 2 other supplemental tests for confirm of HIV infection as per NACO guidelines.

• All patients' reports inderminate should be repeated with a second sample taken 14-28 days. In case the serological results continue to be inderminate the sample should be subject to western blot for confirmation.

Correlate Clinically.

Result rechecked and verified for abnormal cases

*** End Of Report ***





DR. RUTURAJ MANIKLAL KOLHAPURE MD, MICROBIOLOGIST

Note : This report is subject to the terms and conditions overleaf. Partial Reproduction of this report is not Permitted