

# Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

### REPORT

Name : Mrs. BHASWATI CHATTERJEE Sample ID : A0093786
Age/Gender : 33 Years/Female Reg. No : 0312403070051
Referred by : Dr. Nivedita Ashrit MD (Obs/Gyn) SPP Code : SPL-CV-172
Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 07-Mar-2024 07:4

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 07-Mar-2024 07:40 PM
Primary Sample : Whole Blood Received On : 07-Mar-2024 09:50 PM
Sample Tested In : Whole Blood EDTA Reported On : 07-Mar-2024 10:08 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

HAEMATOLOGY				
Test Name	Results	Units	Ref. Range	Method
Complete Blood Picture(CBP)				
Haemoglobin (Hb)	10.0	g/dL	12-15	Cynmeth Method
Haematocrit (HCT)	31.4	%	40-50	Calculated
RBC Count	4.83	10^12/L	4.5-5.5	Cell Impedence
MCV	65	fl	81-101	Calculated
MCH	20.7	pg	27-32	Calculated
MCHC	31.8	g/dL	32.5-34.5	Calculated
RDW-CV	16.6	%	11.6-14.0	Calculated
Platelet Count (PLT)	148	10^9/L	150-410	Cell Impedance
Total WBC Count	10.7	10^9/L	4.0-10.0	Impedance
Differential Leucocyte Count (DC)				
Neutrophils	67	%	40-70	Cell Impedence
Lymphocytes	28	%	20-40	Cell Impedence
Monocytes	04	%	2-10	Microscopy
Eosinophils	01	%	1-6	Microscopy
Basophils	0	%	1-2	Microscopy
Absolute Neutrophils Count	7.17	10^9/L	2.0-7.0	Impedence
Absolute Lymphocyte Count	3	10^9/L	1.0-3.0	Impedence
Absolute Monocyte Count	0.43	10^9/L	0.2-1.0	Calculated
Absolute Eosinophils Count	0.11	10^9/L	0.02-0.5	Calculated
Absolute Basophil ICount	0.00	10^9/L	0.0-0.3	Calculated
Morphology	,		ytic normochromic and few nd Mild Leucocytosis	PAPs Staining







Swarnabala - M DR.SWARNA BALA MD PATHOLOGY



## Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

### REPORT

Name : Mrs. BHASWATI CHATTERJEE Sample ID : A0093785 Age/Gender : 33 Years/Female Reg. No : 0312403070051 Referred by : Dr. Nivedita Ashrit MD (Obs/Gyn) SPP Code : SPL-CV-172 Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 07-Mar-2024 07:40 PM Primary Sample : Whole Blood Received On : 07-Mar-2024 09:50 PM Sample Tested In : Serum Reported On : 07-Mar-2024 10:48 PM

Client Address : Kimtee colony , Gokul Nagar, Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY				
Test Name	Results	Units	Ref. Range	Method
25 - Hydroxy Vitamin D	29.51	ng/mL	<20.0-Deficiency	CLIA
		-	20.0-<30.0-Insufficiency	/
			30.0-100.0-Sufficiency	
			>100.0-Potential Intoxic	ation

#### **Interpretation:**

- 1. Vitamin D helps your body absorb calcium and maintain strong bones throughout your entire life. Your body produces vitamin D when the sun's UV rays contact your skin. Other good sources of the vitamin include fish, eggs, and fortified dairy products. It's also available as a dietary supplement.
- 2. Vitamin D must go through several processes in your body before your body can use it. The first transformation occurs in the liver. Here, your body converts vitamin D to a chemical known as 25-hydroxyvitamin D, also called calcidiol.
- $\bf 3$ . The 25-hydroxy vitamin D test is the best way to monitor vitamin D levels. The amount of 25-hydroxyvitamin D in your blood is a good indication of how much vitamin D your body has. The test can determine if your vitamin D levels are too high or too low.
- **4**. The test is also known as the 25-OH vitamin D test and the calcidiol 25-hydroxycholecalcifoerol test. It can be an important indicator of osteoporosis (bone weakness) and rickets (bone malformation).

#### Those who are at high risk of having low levels of vitamin D include:

1.people who don't get much exposure to the sun

2.older adults

3.people with obesity.

4. dietary deficiency

Increased Levels: Vitamin D Intoxication

Method : CLIA











## Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## REPORT

Name : Mrs. BHASWATI CHATTERJEE

Age/Gender : 33 Years/Female

Referred by : Dr. Nivedita Ashrit MD (Obs/Gyn)
Referring Customer : V CARE MEDICAL DIAGNOSTICS

Primary Sample : Whole Blood

Sample Tested In : Serum

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

Sample ID : A0093785

Reg. No : 0312403070051

SPP Code : SPL-CV-172

Collected On : 07-Mar-2024 07:40 PM

Reported On : 07-Mar-2024 10:48 PM

: 07-Mar-2024 09:50 PM

Report Status Final Report

Received On

### **CLINICAL BIOCHEMISTRY**

Test Name	Results	Units	Ref. Range	Method	
TSH -Thyroid Stimulating Hormone	3.43	μIU/mL	0.35-5.5	CLIA	

#### Pregnancy & Cord Blood

		TSH (Thyroid Stimulating Hormone (μIU/mL)	
First Trimester	: 0.24-2.99		
Second Trimester	: 0.46-2.95		
Third Trimester	: 0.43-2.78		
Cord Blood	: 2.3-13.2		

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

Correlate Clinically.

Result rechecked and verified for abnormal cases

Laboratory is NABL Accredited

\*\*\* End Of Report \*\*\*







DR. VAISHNAVI MD BIOCHEMISTRY