

**REPORT**

Name	: Mrs. T HANISHA	Sample ID	: 24864536
Age/Gender	: 25 Years/Female	Reg. No	: 0312405080028
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 08-May-2024 01:30 PM
Primary Sample	: Whole Blood	Received On	: 08-May-2024 03:47 PM
Sample Tested In	: Serum	Reported On	: 08-May-2024 06:43 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

**CLINICAL BIOCHEMISTRY**

Test Name	Results	Units	Ref. Range	Method
<b>Creatinine -Serum</b>	0.63	mg/dL	0.60-1.10	Sarcosine oxidase

**Interpretation:**

- This test is done to see how well your kidneys are working.Creatinine is a chemical waste product of creatine. Creatine is a chemical made by the body and is used to supply energy mainly to muscles.
- **A higher than normal level may be due to:**  
Renal diseases and insufficiency with decreased glomerular filtration, urinary tract obstruction, reduced renal blood flow including congestive heart failure, shock, and dehydration; rhabdomyolysis can cause elevated serum creatinine.
- **A lower than normal level may be due to:**  
Small stature, debilitation, decreased muscle mass; some complex cases of severe hepatic disease can cause low serum creatinine levels. In advanced liver disease, low creatinine may result from decreased hepatic production of creatinine and inadequate dietary protein as well as reduced muscle mass.

<b>Beta- Human Chorionic Gonodotropin Hormone</b>	3365	mIU/mL	Refer to Interpretation	CLIA
---	------	--------	-------------------------	------

**Interpretation:**

- A quantitative human chorionic gonadotropin (HCG) test measures the specific level of HCG in the blood. HCG is a hormone produced in the body during pregnancy.
- HCG appears in the blood and urine of pregnant women as early as 10 days after conception. Quantitative HCG measurement helps determine the exact age of the fetus. It can also assist in the diagnosis of abnormal pregnancies, such as ectopic pregnancies, molar pregnancies, and possible miscarriages. It is also used as part of a screening test for Down syndrome.
- This test is also done to diagnose abnormal conditions not related to pregnancy that can raise HCG level.

**Non Pregnant Females: < 10.0 mIU/mL**

**Post Menopausal Females: < 10.0 mIU/mL**

**Pregnancy**

Gestational Age and Expected hCG Values (mIU/mL)	Gestational Age and Expected hCG Values (mIU/mL)	Gestational Age and Expected hCG Values (mIU/mL)
0.2-1 weeks: 10-50	1-2 weeks : 50-500	2-3 weeks : 500-10,000
3-4 weeks : 1000-50,000	5-6 weeks : 10,000-100,000	6-8 weeks : 15,000-200,000
2-3 months : 10,000-100,000		

Correlate Clinically.

Laboratory is NABL Accredited

\*\*\* End Of Report \*\*\*



*Dr. Vaishnavi*  
**DR. VAISHNAVI**  
**MD BIOCHEMISTRY**