

# Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

# REPORT

Name : Mrs. SULOCHANA Sample ID : A0286710, A0286709, 248645

Age/Gender : 76 Years/Female Reg. No : 0312405120001

Referred by : Dr. SITHAL KUMAR SPP Code : SPL-CV-172
Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 12-May-2024 07:44 AM

Primary Sample : Whole Blood Received On : 12-May-2024 03:14 PM

Sample Tested In : Plasma-NaF(F), Plasma-NaF(PP), Reported On : 12-May-2024 04:05 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

## **CLINICAL BIOCHEMISTRY**

mg/dL

Test Name Results Units Ref. Range Method

**Glucose Fasting (F) 189**mg/dL
70-100
GOD-POD

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	>= 126	>= 200	>= 6.5	>=200(with symptoms)

Reference: Diabetes care 2018:41(suppl.1):S13-S27

### Glucose Post Prandial (PP) 293

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	>= 126	>= 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2018:41(suppl.1):S13-S27

- Postprandial glucose level is a screening test for Diabetes Mellitus
- If glucose level is >140 mg/dL and <200 mg/dL, then GTT (glucose tolerance test) is advised.</li>
- If level after 2 hours = >200 mg/dL diabetes mellitus is confirmed.
- · Advise HbA1c for further evaluation.

Glycated Hemoglobin (HbA1c) 9.2 % Non Diabetic:< 5.7 HPLC

Pre diabetic: 5.7-6.4 Diabetic:>= 6.5

70-140

Mean Plasma Glucose 217.34 mg/dL Calculated

## Interpretation:

- Glycated hemoglobins (GHb), also called glycohemoglobins, are substances formed when glucose binds to hemoglobin, and occur in amounts proportional to the concentration of serum glucose. Since red blood cells survive an average of 120 days, the measurement of GHb provides an index of a person's average blood glucose concentration (glycemia) during the preceding 2-3 months. Normally, only 4% to 6% of hemoglobin is bound to glucose, while elevated glycohemoglobin levels are seen in diabetes and other hyperglycemic states
- Mean Plasma Glucose(MPG): This Is Mathematical Calculations Where Glycated Hb Can Be Correlated With Daily Mean Plasma Glucose Level

Correlate Clinically.

Result rechecked and verified for abnormal cases

Laboratory is NABL Accredited

\*\*\* End Of Report \*\*\*







DR.VAISHNAVI MD BIOCHEMISTRY

Hexokinase (HK)