



Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## REPORT

Name : Mrs. M MADHURI Sample ID : A0287278

Age/Gender : 71 Years/Female Reg. No : 0312406180002

Referred by : Dr. PADMAVATHI SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 18-Jun-2024 08:12 AM

Primary Sample : Whole Blood Received On : 18-Jun-2024 12:57 PM Sample Tested In : Whole Blood EDTA Reported On : 18-Jun-2024 01:42 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

### **HAEMATOLOGY**

#### **SAGEPATH CARE 1.2**

Test Name	Results	Units	Ref. Range	Method
COMPLETE BLOOD COUNT (CBC)				
Haemoglobin (Hb)	10.5	g/dL	12-15	Cynmeth Method
RBC Count	3.96	10^12/L	4.5-5.5	Cell Impedence
Haematocrit (HCT)	31.8	%	40-50	Calculated
MCV	80	fl	81-101	Calculated
MCH	26.5	pg	27-32	Calculated
MCHC	30.0	g/dL	32.5-34.5	Calculated
RDW-CV	15.7	%	11.6-14.0	Calculated
Platelet Count (PLT)	268	10^9/L	150-410	Cell Impedance
Total WBC Count	9.4	10^9/L	4.0-10.0	Impedance
Neutrophils	64	%	40-70	Cell Impedence
Absolute Neutrophils Count	6.02	10^9/L	2.0-7.0	Impedence
Lymphocytes	30	%	20-40	Cell Impedence
Absolute Lymphocyte Count	2.82	10^9/L	1.0-3.0	Impedence
Monocytes	04	%	2-10	Microscopy
Absolute Monocyte Count	0.38	10^9/L	0.2-1.0	Calculated
Eosinophils	02	%	1-6	Microscopy
Absolute Eosinophils Count	0.19	10^9/L	0.02-0.5	Calculated
Basophils	00	%	1-2	Microscopy
Absolute Basophil ICount	0.00	10^9/L	0.0-0.3	Calculated
<u>Morphology</u>				
WBC	Within Norr	mal Limits		
RBC	Anisocytos	is with Microcyt	tic hypochromic anemia	
Platelets	Adequate.			Microscopy

Result rechecked and verified for abnormal cases

\*\*\* End Of Report \*\*\*

Laboratory is NABL Accredited







Swarnabala - M DR.SWARNA BALA MD PATHOLOGY



Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

### REPORT

Name : Mrs. M MADHURI

Age/Gender : 71 Years/Female Referred by : Dr. PADMAVATHI

Referring Customer: V CARE MEDICAL DIAGNOSTICS

Primary Sample : Whole Blood

Sample Tested In : Whole Blood EDTA

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

Sample ID : A0287278

Reg. No : 0312406180002

SPP Code : SPL-CV-172

Collected On : 18-Jun-2024 08:12 AM

Received On : 18-Jun-2024 12:57 PM

Reported On : 18-Jun-2024 01:46 PM

Report Status : Final Report

### **HAEMATOLOGY**

#### **SAGEPATH CARE 1.2**

Test Name Results Units Ref. Range Method

Erythrocyte Sedimentation Rate (ESR) 44 30 or less Westergren method

Comments: ESR is an acute phase reactant which indicates presence and intensity of an inflammatory process. It is never diagnostic of a specific disease. It is used to monitor the course or response to treatment of certain diseases. Extremely high levels are found in cases of malignancy, hematologic diseases, collagen disorders and renal diseases.









Swarnabala - M DR.SWARNA BALA MD PATHOLOGY



Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## REPORT

Name : Mrs. M MADHURI

Age/Gender : 71 Years/Female Referred by : Dr. PADMAVATHI

Referring Customer : V CARE MEDICAL DIAGNOSTICS

Primary Sample : Whole Blood Sample Tested In : Plasma-NaF(F)

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

Sample ID : A0287276

Reg. No : 0312406180002

SPP Code : SPL-CV-172

Collected On : 18-Jun-2024 08:12 AM

Received On : 18-Jun-2024 12:57 PM

Reported On : 18-Jun-2024 01:37 PM Report Status : Final Report

### **CLINICAL BIOCHEMISTRY**

#### **SAGEPATH CARE 1.2**

Test Name Results Units Ref. Range Method

Glucose Fasting (F) 83 mg/dL 70-100 Hexokinase

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	>= 126	>= 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2018:41(suppl.1):S13-S27

Result rechecked and verified for abnormal cases

\*\*\* End Of Report \*\*\*

Laboratory is NABL Accredited

Excellence In Health Care









Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## REPORT

Name: Mrs. M MADHURISample ID: A0287278Age/Gender: 71 Years/FemaleReg. No: 0312406180002Referred by: Dr. PADMAVATHISPP Code: SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 18-Jun-2024 08:12 AM
Primary Sample : Whole Blood Received On : 18-Jun-2024 12:57 PM
Sample Tested In : Whole Blood EDTA Reported On : 18-Jun-2024 01:37 PM

Client Address : Kimtee colony , Gokul Nagar, Tarnaka Report Status : Final Report

#### **CLINICAL BIOCHEMISTRY**

#### **SAGEPATH CARE 1.2**

OAGELATII OARE 112					
Test Name	Results	Units	Ref. Range	Method	
Glycated Hemoglobin (HbA1c)	6.4	%	Non Diabetic: < 5.7 Pre diabetic: 5.7-6.4	HPLC	
Mean Plasma Glucose	136.98	mg/dL	Diabetic:>= 6.5	Calculated	

Glycated hemoglobins (GHb), also called glycohemoglobins, are substances formed when glucose binds to hemoglobin, and occur in amounts proportional to the concentration of serum glucose. Since red blood cells survive an average of 120 days, the measurement of GHb provides an index of a person's average blood glucose concentration (glycemia) during the preceding 2-3 months. Normally, only 4% to 6% of hemoglobin is bound to glucose, while elevated glycohemoglobin levels are seen in diabetes and other hyperglycemic states Mean Plasma Glucose(MPG):This Is Mathematical Calculations Where Glycated Hb Can Be Correlated With Daily Mean Plasma Glucose Level

NOTE: The above Given Risk Level Interpretation is not age specific and is an information resource only and is not to be used or relied on for any diagnostic or treatment purposes and should not be used as a substitute for professional diagnosis and treatment. Kindly Correlate clinically.

#### INTERPRETATION

### Method: Analyzer Fully automated HPLC platform.

Average Blood Glucose(eAG) (mg/dL)	Level of Control	Hemoglobin A10 (%)
421		14%
386	A	13%
350	L	12%
314	E	11%
279	R	10%
243	T	9%
208		8%
172	POOR	7%
136	GOOD	6%
101	EXCELLENT	5%

HbA1c values of 5.0- 6.5 percent indicate good control or an increased risk for developing diabetes mellitus. HbA1c values greater than 6.5 percent are diagnostic of diabetes mellitus. Diagnosis should be confirmed by repeating the HbA1c test.

NOTE: Hb F higher than 10 percent of total Hb may yield falsely low results. Conditions that shorten red cell survival, such as the presence of unstable hemoglobins like Hb SS, Hb CC, and Hb SC, or other causes of hemolytic anemia may yield falsely low results. Iron deficiency anemia may yield falsely high results.

\*\*\* End Of Report \*\*\*

Laboratory is NABL Accredited









Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## REPORT

Name : Mrs. M MADHURI

Age/Gender : 71 Years/Female Referred by : Dr. PADMAVATHI

Referring Customer : V CARE MEDICAL DIAGNOSTICS

Primary Sample : Whole Blood

Sample Tested In : Serum

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

Sample ID : A0287277

Reg. No : 0312406180002

SPP Code : SPL-CV-172

Collected On : 18-Jun-2024 08:12 AM

Received On : 18-Jun-2024 12:57 PM

Reported On : 18-Jun-2024 03:18 PM

Report Status : Final Report

### **CLINICAL BIOCHEMISTRY**

#### **SAGEPATH CARE 1.2**

rest Name nesults offits net. Name method	Test Name Results Units	Ref. Range Method
---	-------------------------	-------------------

Calcium 8.7 mg/dL 8.5-10.1 Arsenazo

#### Comments:

- Calcium in the body is found mainly in the bones (approximately 99%). In serum, Calcium exists in a
  free ionised form and in bound form (with Albumin). Hence, a decrease in Albumin causes lower
  Calcium levels and vice-versa.
- Calcium levels in serum depend on the Parathyroid Hormone.
- Increased Calcium levels are found in Bone tumors, Hyperparathyroidism. decreased levels are found in Hypoparathyroidism, renal failure, Rickets.

\*\*\* End Of Report \*\*\*

Laboratory is NABL Accredited

Excellence In Health Care









Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## **REPORT**

 Name
 : Mrs. M MADHURI
 Sample ID
 : A0287277

 Age/Gender
 : 71 Years/Female
 Reg. No
 : 0312406180002

Referred by : Dr. PADMAVATHI SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 18-Jun-2024 08:12 AM
Primary Sample : Whole Blood Received On : 18-Jun-2024 12:57 PM
Sample Tested In : Serum Reported On : 18-Jun-2024 03:18 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

### **CLINICAL BIOCHEMISTRY**

#### **SAGEPATH CARE 1.2**

Test Name	Results	Units	Ref. Range	Method
Lipid Profile				
Cholesterol Total	184	mg/dL	< 200	CHOD-POD
Triglycerides-TGL	173	mg/dL	< 150	GPO-POD
Cholesterol-HDL	47	mg/dL	40-60	Direct
Cholesterol-LDL	102.4	mg/dL	< 100	Calculated
Cholesterol- VLDL	34.6	mg/dL	7-35	Calculated
Non HDL Cholesterol	137	mg/dL	< 130	Calculated
Cholesterol Total /HDL Ratio	3.91	%	0-4.0	Calculated
HDL / LDL Ratio	0.46			
LDL/HDL Ratio	2.18	%	0-3.5	Calculated

The National Cholesterol Education program's third Adult Treatment Panel (ATPIII) has issued its recommendations on evaluating and treating lipid discorders for primary and secondary.

NCEP Recommendations	Cholesterol Total in (mg/dL)	Trialveerides	HDL Cholesterol (mg/dL)	LDL Cholesterol	Non HDL Cholesterol in (mg/dL)
( )ntimai	Adult: < 200 Children: < 170	< 150	40-59	Adult:<100 Children: <110	<130
Above Optimal				100-129	130 - 159
Borderline High	Adult: 200-239 Children:171-199	150-199		Adult: 130-159 Children: 111-129	160 - 189
High	Adult:>or=240 Children:>or=200	200-499	≥ 60	Adult:160-189 Children:>or=130	190 - 219
Very High		>or=500		Adult: >or=190 	>=220

Note: LDL cholesterol cannot be calculated if triglyceride is >400 mg/dL (Friedewald's formula). Calculated values not provided for LDL and VLDL

Result rechecked and verified for abnormal cases

\*\*\* End Of Report \*\*\*

Laboratory is NABL Accredited









Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## REPORT

Name : Mrs. M MADHURI Sample ID : A0287277

Age/Gender : 71 Years/Female Reg. No : 0312406180002 Referred by : Dr. PADMAVATHI SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 18-Jun-2024 08:12 AM
Primary Sample : Whole Blood Received On : 18-Jun-2024 12:57 PM

Sample Tested In : Serum Reported On : 18-Jun-2024 03:18 PM

Client Address : Kimtee colony , Gokul Nagar, Tarnaka Report Status : Final Report

### **CLINICAL BIOCHEMISTRY**

#### **SAGEPATH CARE 1.2**

Test Name	Results	Units	Ref. Range	Method
Kidney Profile-KFT				
Creatinine -Serum	0.64	mg/dL	0.55-1.02	Sarcosine oxidase
Urea-Serum	20.9	mg/dL	17.1-49.2	Glutamate dehydrogenase+Calculation
Blood Urea Nitrogen (BUN)	9.77	mg/dL	8.0-23.0	Calculated
BUN / Creatinine Ratio	15.27		6 - 22	
Uric Acid	5.9	mg/dL	2.6-6.0	Uricase
Sodium	138	mmol/L	136-145	ISE Direct
Potassium	3.7	mmol/L	3.5-5.1	ISE Direct
Chloride	102	mmol/L	98-108	ISE Direct

#### Interpretation:

• The kidneys, located in the retroperitoneal space in the abdomen, are vital for patient health. They process several hundred liters of fluid a day and remove around two liters of waste products from the bloodstream. The volume of fluid that passes though the kidneys each minute is closely linked to cardiac output. The kidneys maintain the body's balance of water and concentration of minerals such as sodium, potassium, and phosphorus in blood and remove waste by-products from the blood after digestion, muscle activity and exposure to chemicals or medications. They also produce renin which helps regulate blood pressure, produce erythropoietin which stimulates red blood cell production, and produce an active form of vitamin D, needed for bone health.

\*\*\* End Of Report \*\*\*

Laboratory is NABL Accredited











**Test Na** 



# Sagepath Labs Pvt. Ltd.

Lab Address: - # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

## REPORT

Name : Mrs. M MADHURI Sample ID : A0287277 Age/Gender : 71 Years/Female Reg. No : 0312406180002 SPP Code Referred by : Dr. PADMAVATHI : SPL-CV-172 Referring Customer: V CARE MEDICAL DIAGNOSTICS Collected On : 18-Jun-2024 08:12 AM Primary Sample : Whole Blood : 18-Jun-2024 12:57 PM Received On Sample Tested In : Serum Reported On : 18-Jun-2024 03:18 PM Client Address

: Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

## **CLINICAL BIOCHEMISTRY SAGEPATH CARE 1.2**

ame	Results	Units	Ref. Range	Method

Liver Function Test (LFT)				
Bilirubin(Total)	0.9	mg/dL	0.2-1.2	Diazo
Bilirubin (Direct)	0.2	mg/dL	0.0 - 0.2	Diazo
Bilirubin (Indirect)	0.7	mg/dL	0.2-1.0	Calculated
Aspartate Aminotransferase (AST/SGOT)	27	U/L	5-48	IFCC with out (P-5-P)
Alanine Aminotransferase (ALT/SGPT)	17	U/L	0-55	IFCC with out (P-5-P)
Alkaline Phosphatase(ALP)	101	U/L	30-120	Kinetic PNPP-AMP
Gamma Glutamyl Transpeptidase (GGTP)	17	U/L	5-55	IFCC
Protein - Total	6.4	g/dL	6.4-8.2	Biuret
Albumin	3.4	g/dL	3.4-5.0	Bromocresol Green (BCG)
Globulin	3	g/dL	2.0-4.2	Calculated
A:G Ratio	1.13	%	0.8-2.0	Calculated
SGOT/SGPT Ratio	1.59			

Alanine Aminotransferase(ALT) is an enzyme found in liver and kidneys cells. ALT helps create energy for liver cells. Damaged liver cells release ALT into the bloodstream, which can elevate ALT levels in the blood.

Aspartate Aminotransferase (AST) is an enzyme in the liver and muscles that helps metabolizes amino acids. Similarly to ALT, elevated AST levels may be a sign of liver damage or liver

Alkaline phosphate (ALP) is an enzyme present in the blood. ALP contributes to numerous vital bodily functions, such as supplying nutrients to the liver, promoting bone growth, and

Gamma-glutamyl Transpeptidase (GGTP) is an enzyme that occurs primarily in the liver, but it is also present in the kidneys, pancreas, gallbladder, and spleen. Higher than normal concentrations of GGTP in the blood may indicate alcohol-related liver damage. Elevated GGTP levels can also increase the risk of developing certain types of cancer.

Bilirubin is a waste product that forms when the liver breaks down red blood cells. Bilirubin exits the body as bile in stool. High levels of bilirubin can cause jaundice - a condition in which the skin and whites of the eves turn vellow- and may indicate liver damage.

Albumin is a protein that the liver produces. The liver releases albumin into the bloodstream, where it helps fight infections and transport vitamins, hormones, and enzymes throughout the body. Liver damage can cause abnormally low albumin levels.

\*\*\* End Of Report \*\*\*

Laboratory is NABL Accredited









Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

#### REPOR1

Name : Mrs. M MADHURI Sample ID : A0287277

Age/Gender : 71 Years/Female Reg. No : 0312406180002

Referred by : Dr. PADMAVATHI SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 18-Jun-2024 08:12 AM Primary Sample : Whole Blood Received On : 18-Jun-2024 12:57 PM

Sample Tested In : Serum Reported On : 18-Jun-2024 03:18 PM

Client Address : Kimtee colony , Gokul Nagar, Tarnaka Report Status : Final Report

#### **CLINICAL BIOCHEMISTRY**

#### **SAGEPATH CARE 1.2**

lest Name	Results	Units	Ref. Range	Method	
Thyroid Profile-I(TFT)					
T3 (Triiodothyronine)	132.65	ng/dL	40-181	CLIA	
T4 (Thyroxine)	9.8	μg/dL	3.2-12.6	CLIA	
TSH -Thyroid Stimulating Hormone	4.60	μIU/mL	0.35-5.5	CLIA	

#### Pregnancy & Cord Blood

T3 (Triiodothyronine):		T4 (Thyroxine)	TSH (Thyroid Stimulating Hormone)
First Trimester	: 81-190 ng/dL	15 to 40 weeks:9.1-14.0 μg/dL	First Trimester : 0.24-2.99 µIU/mL
Second&Third Trimeste	er :100-260 ng/dL		Second Trimester: 0.46-2.95 µIU/mL
			Third Trimester : 0.43-2.78 µIU/mL
Cord Blood: 30-70 ng/d	dL/	Cord Blood: 7.4-13.0 µg/dL	Cord Blood: : 2.3-13.2 µIU/mL

#### **Interpretation:**

- Thyroid gland is a butterfly-shaped endocrine gland that is normally located in the lower front of the neck. The thyroid's job is to make thyroid hormones, which are secreted into the blood and then carried to every tissue in the body. Thyroid hormones help the body use energy, stay warm and keep the brain, heart, muscles, and other organs working as they should.
- Thyroid produces two major hormones: triiodothyronine (T3) and thyroxine (T4). If thyroid gland doesn't produce enough of these hormones, you may experience symptoms such as weight gain, lack of energy, and depression. This condition is called hypothyroidism.
- Thyroid gland produces too many hormones, you may experience weight loss, high levels of anxiety, tremors, and a sense of being on a high. This is called hyperthyroidism.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4.
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.











Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

### REPORT

Name : Mrs. M MADHURI Sample ID : A0287277

Age/Gender : 71 Years/Female Reg. No : 0312406180002 Referred by : Dr. PADMAVATHI SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 18-Jun-2024 08:12 AM

Primary Sample : Whole Blood Received On : 18-Jun-2024 12:57 PM Sample Tested In : Serum Reported On : 18-Jun-2024 03:18 PM

Client Address : Kimtee colony , Gokul Nagar, Tarnaka Report Status : Final Report

### **CLINICAL BIOCHEMISTRY**

#### **SAGEPATH CARE 1.2**

Test Name	Results	Units	Ref. Range	Method
Iron Profile-I				
Iron(Fe)	79	μg/dL	50-170	Ferene
Total Iron Binding Capacity (TIBC)	374	μg/dL	250-450	Ferene
Transferrin	261.54	mg/dL	250-380	Calculated
Iron Saturation((% Transferrin Saturation)	21.12	%	15-50	Calculated
Unsaturated Iron Binding Capacity (UIBC)	295	ug/dL	110-370	FerroZine

#### Interpretation:

- Serum transferrin (and TIBC) high, serum iron low, saturation low. Usual causes of depleted iron stores include blood loss, inadequate dietary iron. RBCs in moderately severe iron deficiency are hypochromic and microcytic. Stainable marrow iron is absent. Serum ferritin decrease is the earliest indicator of iron deficiency if inflammation is absent.
- Anemia of chronic disease: Serum transferrin (and TIBC) low to normal, serum iron low, saturation low or normal. Transferrin decreases with many inflammatory diseases. With chronic disease there is a block in movement to and utilization of iron by marrow. This leads to low serum iron and decreased erythropoiesis. Examples include acute and chronic infections, malignancy and renal failure.
- Sideroblastic Anemia: Serum transferrin (and TIBC) normal to low, serum iron normal to high, saturation high.
- Hemolytic Anemia: Serum transferrin (and TIBC) normal to low, serum iron high, saturation high.
- Hemochromatosis: Serum transferrin (and TIBC) slightly low, serum iron high, saturation very high
- Protein depletion: Serum transferrin (and TIBC) may be low, serum iron normal or low (if patient also is iron deficient). This may occur as a result of malnutrition, liver disease, renal disease.
- Liver disease: Serum transferrin variable; with acute viral hepatitis, high along with serum iron and ferritin. With chronic liver disease (eg, cirrhosis), transferrin may be low. Patients who have cirrhosis and portacaval shunting have saturated TIBC/transferrin as well as high ferritin.

Correlate Clinically.

Laboratory is NABL Accredited

\*\*\* End Of Report \*\*\*





