

REPORT

Name	: Mr. V AROGYA REDDY	Sample ID	: A0590410
Age/Gender	: 53 Years/Male	Reg. No	: 0312408020006
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 02-Aug-2024 09:42 AM
Primary Sample	: Whole Blood	Received On	: 02-Aug-2024 01:55 PM
Sample Tested In	: Serum	Reported On	: 02-Aug-2024 05:31 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Ref. Range	Method
Uric Acid	4.4	mg/dL	3.5-7.2	Uricase

Interpretation:

- Uric acid is a chemical created when the body breaks down substances called purines. Purines are normally produced in the body and are also found in some foods and drinks. Foods with high content of purines include liver, anchovies, mackerel, dried beans and peas, and beer. Most uric acid dissolves in blood and travels to the kidneys. From there, it passes out in urine. If your body produces too much uric acid or does not remove enough of it, you can get sick. A high level of uric acid in the blood is called hyperuricemia. This test checks to see how much uric acid you have in your blood. Investigation and monitoring of inflammatory arthritis pain, particularly in big toe (gout)
- Useful in the investigation of kidney stones
- Aid in diagnosis, treatment, and monitoring of renal failure/disease
- Monitor patients receiving cytotoxic drugs (high nucleic acid turnover)
- Monitor diseases with nucleic acid metabolism and turnover (eg, leukemia, lymphoma, polycythemia)



Creatinine -Serum	0.8	mg/dL	0.70-1.30	Jaffes Kinetic
-------------------	-----	-------	-----------	----------------

Interpretation:

- This test is done to see how well your kidneys are working. Creatinine is a chemical waste product of creatine. Creatine is a chemical made by the body and is used to supply energy mainly to muscles.
- **A higher than normal level may be due to:**
- Renal diseases and insufficiency with decreased glomerular filtration, urinary tract obstruction, reduced renal blood flow including congestive heart failure, shock, and dehydration; rhabdomyolysis can cause elevated serum creatinine.
- **A lower than normal level may be due to:**
- Small stature, debilitation, decreased muscle mass; some complex cases of severe hepatic disease can cause low serum creatinine levels. In advanced liver disease, low creatinine may result from decreased hepatic production of creatinine and inadequate dietary protein as well as reduced muscle mass.



Dr. Vaishnavi
DR. VAISHNAVI
MD BIOCHEMISTRY

REPORT

Name	: Mr. V AROGYA REDDY	Sample ID	: A0590350
Age/Gender	: 53 Years/Male	Reg. No	: 0312408020006
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 02-Aug-2024 09:42 AM
Primary Sample	:	Received On	: 02-Aug-2024 01:47 PM
Sample Tested In	: Urine	Reported On	: 02-Aug-2024 04:14 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

CLINICAL PATHOLOGY

Test Name	Results	Units	Ref. Range	Method
Complete Urine Analysis (CUE)				
Physical Examination				
Colour	Pale Yellow		Straw to light amber	
Appearance	Clear		Clear	
Chemical Examination				
Glucose	Negative		Negative	Strip Reflectance
Protein	Absent		Negative	Strip Reflectance
Bilirubin (Bile)	Negative		Negative	Strip Reflectance
Urobilinogen	Negative		Negative	Ehrlichs reagent
Ketone Bodies	Negative		Negative	Strip Reflectance
Specific Gravity	1.010		1.000 - 1.030	Strip Reflectance
Blood	Negative		Negative	Strip Reflectance
Reaction (pH)	6.0		5.0 - 8.5	Reagent Strip Reflectance
Nitrites	Negative		Negative	Strip Reflectance
Leukocyte esterase	Negative		Negative	Reagent Strip Reflectance
Microscopic Examination (Microscopy)				
PUS(WBC) Cells	02-03	/hpf	00-05	Microscopy
R.B.C.	Nil	/hpf	Nil	Microscopic
Epithelial Cells	01-02	/hpf	00-05	Microscopic
Casts	Absent		Absent	Microscopic
Crystals	Absent		Absent	Microscopic
Bacteria	Nil		Nil	
Budding Yeast Cells	Nil		Absent	Microscopy

Comments :Urine analysis is one of the most useful laboratory tests as it identifies a wide range of medical conditions including renal damage, urinary tract infections,diabetes, hypertension and drug toxicity.

Correlate Clinically.

Laboratory is NABL Accredited

*** End Of Report ***



Swannabala - M
DR.SWARNA BALA
MD PATHOLOGY