



Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

Hexokinase

REPORT

 Name
 : Mr. GIRIDAR RAO
 Sample ID
 : A0590514, A0590508

 Age/Gender
 : 58 Years/Male
 Reg. No
 : 0312408040010

 Referred by
 : Dr. SELF
 SPP Code
 : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 04-Aug-2024 08:08 AM Primary Sample : Whole Blood Received On : 04-Aug-2024 06:20 PM

Sample Tested In : Plasma-NaF(F), Plasma-NaF(PP) Reported On : 04-Aug-2024 07:28 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY

GLUCOSE POST PRANDIAL (PP)

Test Name Results Units Ref. Range Method

Glucose Fasting (F) 111 mg/dL

Reference: Diabetes care 2018:41(suppl.1):S13-S27

Glucose Post Prandial (PP)

171 mg/dL 70-140 Hexokinase (HK)

70-100

]	Interpretation of	Plasma	Glucose	based	on AI	DA g	guidelines	2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	>= 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2018:41(suppl.1):S13-S27

- Postprandial glucose level is a screening test for Diabetes Mellitus
- If glucose level is >140 mg/dL and <200 mg/dL, then GTT (glucose tolerance test) is advised.
- If level after 2 hours = >200 mg/dL diabetes mellitus is confirmed.
- Advise HbA1c for further evaluation.

Result rechecked and verified for abnormal cases

*** End Of Report ***

Laboratory is NABL Accredited







DR.VAISHNAVI MD BIOCHEMISTRY



Sagepath Labs Pvt. Ltd.

Lab Address: - # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

: A0590511

REPORT

Sample ID

Name : Mr. GIRIDAR RAO

Age/Gender : 58 Years/Male Reg. No : 0312408040010

Referred by : Dr. SELF SPP Code : SPL-CV-172

Referring Customer: V CARE MEDICAL DIAGNOSTICS

Collected On : 04-Aug-2024 08:08 AM Primary Sample : Whole Blood : 04-Aug-2024 06:20 PM Received On

Sample Tested In : Whole Blood EDTA Reported On : 04-Aug-2024 06:55 PM Report Status : Final Report

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

CLINICAL BIOCHEMISTRY

CEINICAE BIOCHEMISTR I					
Test Name	Results	Units	Ref. Range	Method	
Glycated Hemoglobin (HbA1c)	11.0	%	Non Diabetic:< 5.7 Pre diabetic: 5.7-6.4 Diabetic:>= 6.5	HPLC	
Mean Plasma Glucose	269	mg/dL		Calculated	

Glycated hemoglobins (GHb), also called glycohemoglobins, are substances formed when glucose binds to hemoglobin, and occur in amounts proportional to the concentration of serum glucose. Since red blood cells survive an average of 120 days, the measurement of GHb provides an index of a person's average blood glucose concentration (glycemia) during the preceding 2-3 months. Normally, only 4% to 6% of hemoglobin is bound to glucose, while elevated glycohemoglobin levels are seen in diabetes and other hyperglycemic states Mean Plasma Glucose(MPG): This Is Mathematical Calculations Where Glycated Hb Can Be Correlated With Daily Mean Plasma Glucose Level

NOTE: The above Given Risk Level Interpretation is not age specific and is an information resource only and is not to be used or relied on for any diagnostic or treatment purposes and should not be used as a substitute for professional diagnosis and treatment. Kindly Correlate clinically.

INTERPRETATION

Method: Analyzer Fully automated HPLC platform.

Average Blood Glucose(eAG) (mg/dL)	Level of Control	Hemoglobin A10 (%)	
421		14%	
386	_ A _	13%	
350	L	12%	
314	E	11%	
279	R	10%	
243	Т	9%	
208		8%	
172	POOR	7%	
136	GOOD	6%	
101	EXCELLENT	5%	

HbA1c values of 5.0- 6.5 percent indicate good control or an increased risk for developing diabetes mellitus. HbA1c values greater than 6.5 percent are diagnostic of diabetes mellitus. Diagnosis should be confirmed by repeating the HbA1c test.

NOTE: Hb F higher than 10 percent of total Hb may yield falsely low results. Conditions that shorten red cell survival, such as the presence of unstable hemoglobins like Hb SS, Hb CC, and Hb SC, or other causes of hemolytic anemia may yield falsely low results. Iron deficiency anemia may yield falsely high results.

Correlate Clinically.

Result rechecked and verified for abnormal cases

Laboratory is NABL Accredited

*** End Of Report ***





